

Pubblicazioni 2004-2016

- Sella, L., Castiglioni, C., Paccanaro, M.C., Janni, M., Schäfer, W., D'Ovidio, R., Favaron, F., 2016. Involvement of Fungal Pectin Methyltransferase Activity in the Interaction Between *Fusarium graminearum* and Wheat Mol. Plant. Microbe. Interact. 29, 258–267.
- Sella, L., Gazzetti, K., Castiglioni, C., Schäfer, W., D'Ovidio, R., Favaron, F., 2016. The *Fusarium graminearum* Xyr1 transcription factor regulates xylanase expression but is not essential for fungal virulence. Plant Pathol. n/a–n/a. doi:10.1111/ppa.12456
- Tundo, S., Moschetti, I., Faoro, F., Lafond, M., Giardina, T., Favaron, F., Sella, L., D'Ovidio, R., 2015. *Fusarium graminearum* produces different xylanases causing host cell death that is prevented by the xylanase inhibitors XIP-I and TAXI-III in wheat. Plant Sci. 240, 161–169. doi:10.1016/j.plantsci.2015.09.002
- Moschetti, I., Faoro, F., Moro, S., Sabbadin, D., Sella, L., Favaron, F., D'Ovidio, R., 2015. The xylanase inhibitor TAXI-III counteracts the necrotic activity of a *Fusarium graminearum* xylanase in vitro and in durum wheat transgenic plants. Mol. Plant Pathol. 16, 583–592. doi:10.1111/mpp.12215
- Kalunke, R.M., Cenci, A., Volpi, C., O'Sullivan, D.M., Sella, L., Favaron, F., Cervone, F., De Lorenzo, G., D'Ovidio, R., 2014. The pgip family in soybean and three other legume species: evidence for a birth-and-death model of evolution. BMC Plant Biol. 14, 189. doi:10.1186/s12870-014-0189-3
- Sella, L., Gazzetti, K.G., Castiglioni, C., Schäfer, W., Favaron, F., 2014. *Fusarium graminearum* possesses virulence factors common to Fusarium head blight of wheat and seedling rot of soybean, but differing in their impact on disease severity. Phytopathology 104, 1–30. doi:10.1094/PHTO-12-13-0355-R
- Moschetti, I., Tundo, S., Janni, M., Sella, L., Gazzetti, K., Tauzin, A., Giardina, T., Masci, S., Favaron, F., D'Ovidio, R., 2013. Constitutive expression of the xylanase inhibitor TAXI-III delays Fusarium head blight symptoms in durum wheat transgenic plants. Mol. Plant. Microbe. Interact. 26, 1464–72. doi:10.1094/MPMI-04-13-0121-R
- Sella, L., Gazzetti, K., Faoro, F., Odorizzi, S., D'Ovidio, R., Schäfer, W., Favaron, F., 2013. A *Fusarium graminearum* xylanase expressed during wheat infection is a necrotizing factor but is not essential for virulence. Plant Physiol. Biochem. 64, 1–10. doi:10.1016/j.plaphy.2012.12.008
- Volpi, C., Raiola, A., Janni, M., Gordon, A., O'Sullivan, D.M., Favaron, F., D'Ovidio, R., 2013. *Claviceps purpurea* expressing polygalacturonases escaping PGIP inhibition fully infects PVPGP2 wheat transgenic plants but its infection is delayed in wheat transgenic plants with increased level of pectin methyl esterification. Plant Physiol. Biochem. 73, 294–301. doi:10.1016/j.plaphy.2013.10.011
- Ferrari, S., Sella, L., Janni, M., De Lorenzo, G., Favaron, F., D'Ovidio, R., 2012. Transgenic expression of polygalacturonase-inhibiting proteins in Arabidopsis and wheat increases resistance to the flower pathogen *Fusarium graminearum*. Plant Biol. 14, 31–38. doi:10.1111/j.1438-8677.2011.00449.x
- Kalunke, R.M., Janni, M., Sella, L., David, P., Geffroy, V., Favaron, F., D'Ovidio, R., 2011. Transcript analysis of the bean polygalacturonase inhibiting protein gene family reveals that PVPGP2 is expressed in the whole plant and is strongly induced by pathogen infection. J. Plant Pathol. 93, 141–148.
- Volpi, C., Janni, M., Lionetti, V., Bellincampi, D., Favaron, F., D'Ovidio, R., 2011. The ectopic expression of a pectin methyl esterase inhibitor increases pectin methyl esterification and limits fungal diseases in wheat. Mol. Plant. Microbe. Interact. 24, 1012–1019. doi:10.1094/MPMI-01-11-0021
- Sella, L., Cosmi, T., Giacomello, F., Saccardi, A., Favaron, 2010. First report of *Fusarium oxysporum* on *Dipladenia* sp. in Italy. J. J. Plant Pathol. 92, 543.
- Favaron, F., Lucchetta, M., Odorizzi, S., Pais da Cunha, A.T., Sella, L., 2009. The role of grape polyphenols on

trans-resveratrol activity against *Botrytis cinerea* and of fungal laccase on the solubility of putative grape proteins. *J. Plant Pathol.* 91, 579–588. doi:10.4454/jpp.v91i3.549

- Tomassini, A., Sella, L., Raiola, A., D'Ovidio, R., Favaron, F., 2009. Characterization and expression of *Fusarium graminearum* endo-polygalacturonases in vitro and during wheat infection. *Plant Pathol.* 58, 556–564. doi:10.1111/j.1365-3059.2008.02019.x
- Janni, M., Sella, L., Favaron, F., Blechl, A.E., De Lorenzo, G., D'Ovidio, R., 2008. The expression of a bean PGIP in transgenic wheat confers increased resistance to the fungal pathogen *Bipolaris sorokiniana*. *Mol. Plant. Microbe. Interact.* 21, 171–177. doi:10.1094/MPMI-21-2-0171
- Raiola, A., Sella, L., Castiglioni, C., Balmas, V., Favaron, F., 2008. A single amino acid substitution in highly similar endo-PGs from *Fusarium verticillioides* and related *Fusarium* species affects PGIP inhibition. *Fungal Genet. Biol.* 45, 776–789. doi:10.1016/j.fgb.2007.11.003
- Lionetti, V., Raiola, A., Camardella, L., Giovane, A., Obel, N., Pauly, M., Favaron, F., Cervone, F., Bellincampi, D., 2007. Overexpression of pectin methylesterase inhibitors in *Arabidopsis* restricts fungal infection by *Botrytis cinerea*. *Plant Physiol.* 143, 1871–1880. doi:10.1104/pp.106.090803
- D'Ovidio, R., Roberti, S., Di Giovanni, M., Capodicasa, C., Melaragni, M., Sella, L., Tosi, P., Favaron, F., 2006. The characterization of the soybean polygalacturonase-inhibiting proteins (Pgip) gene family reveals that a single member is responsible for the activity detected in soybean tissues. *Planta* 224, 633–645. doi:10.1007/s00425-006-0235-y
- Sella, L., Tomassini, A., D'Ovidio, R., Favaron, F., 2005. Expression of two *Sclerotinia sclerotiorum* endo-PG genes correlates with endo-polygalacturonase activity during *Glycine max* colonization. *J. Plant Pathol.* 87, 199–205.
- Zuppini, A., Navazio, L., Sella, L., Castiglioni, C., Favaron, F., Mariani, P., 2005. An endopolygalacturonase from *Sclerotinia sclerotiorum* induces calcium-mediated signaling and programmed cell death in soybean cells. *Mol. Plant. Microbe. Interact.* 18, 849–855. doi:10.1094/MPMI-18-0849
- Favaron, F., Sella, L., D'Ovidio, R., 2004. Relationships among endo-polygalacturonase, oxalate, pH, and plant polygalacturonase-inhibiting protein (PGIP) in the interaction between *Sclerotinia sclerotiorum* and soybean. *Mol. Plant. Microbe. Interact.* 17, 1402–1409. doi:10.1094/MPMI.2004.17.12.1402
- Sella, L., Castiglioni, C., Roberti, S., D'Ovidio, R., Favaron, F., 2004. An endo-polygalacturonase (PG) of *Fusarium moniliforme* escaping inhibition by plant polygalacturonase-inhibiting proteins (PGIPs) provides new insights into the PG-PGIP interaction. *FEMS Microbiol. Lett.* 240, 117–124. doi:10.1016/j.femsle.2004.09.019
- Zuppini, A., Baldan, B., Millionsi, R., Favaron, F., Navazio, L., Mariani, P., 2004. Chitosan induces Ca²⁺-mediated programmed cell death in soybean cells. *New Phytol.* 161, 557–568. doi:10.1046/j.1469-8137.2003.00969.x